

## DIRECTIONS FOR USE

B. Braun Melsungen AG · 34209 Melsungen, Germany

# Nutriflex Lipid plus emulsion for infusion

## 1. NAME OF THE MEDICINAL PRODUCT

Nutriflex Lipid plus emulsion for infusion

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

The ready-for-use emulsion for intravenous infusion contains after mixing the chamber contents:

from the upper, left-hand chamber (glucose solution)				
Glucose monohydrate	132.0 g	165.0 g	247.5 g	330.0 g
equivalent to glucose	120.0 g	150.0 g	225.0 g	300.0 g
Sodium dihydrogen phosphate dihydrate	1.872 g	2.340 g	3.510 g	4.680 g
Zinc acetate dihydrate	5.26 mg	6.58 mg	9.87 mg	13.16 mg

from the upper, right-hand chamber (fat emulsion)				
Soya-bean oil, refined	20.0 g	25.0 g	37.5 g	50.0 g
Medium-chain triglycerides	20.0 g	25.0 g	37.5 g	50.0 g

from the lower chamber (amino acid solution)				
Isoleucine	2.26 g	2.82 g	4.23 g	5.64 g
Leucine	3.01 g	3.76 g	5.64 g	7.52 g
Lysine hydrochloride	2.73 g	3.41 g	5.12 g	6.82 g
equivalent to lysine	2.18 g	2.73 g	4.10 g	5.46 g
Methionine	1.88 g	2.35 g	3.53 g	4.70 g
Phenylalanine	3.37 g	4.21 g	6.32 g	8.42 g
Threonine	1.74 g	2.18 g	3.27 g	4.36 g
Tryptophan	0.54 g	0.68 g	1.02 g	1.36 g
Valine	2.50 g	3.12 g	4.68 g	6.24 g
Arginine	2.59 g	3.24 g	4.86 g	6.48 g
Histidine hydrochloride monohydrate	1.62 g	2.03 g	3.05 g	4.06 g
equivalent to histidine	1.20 g	1.50 g	2.25 g	3.00 g
Alanine	4.66 g	5.82 g	8.73 g	11.64 g
Aspartic acid	1.44 g	1.80 g	2.70 g	3.60 g
Glutamic acid	3.37 g	4.21 g	6.32 g	8.42 g
Glycine	1.58 g	1.98 g	2.97 g	3.96 g
Proline	3.26 g	4.08 g	6.12 g	8.16 g
Serine	2.88 g	3.60 g	5.40 g	7.20 g
Sodium hydroxide	0.781 g	0.976 g	1.464 g	1.952 g
Sodium chloride	0.402 g	0.503 g	0.755 g	1.006 g
Sodium acetate trihydrate	0.222 g	0.277 g	0.416 g	0.554 g
Potassium acetate	2.747 g	3.434 g	5.151 g	6.868 g

tetrahydrate	0.686 g	0.858 g	1.287 g	1.716 g
Calcium chloride dihydrate	0.470 g	0.588 g	0.882 g	1.176 g

	in 1000 ml	in 1250 ml	in 1875 ml	in 2500 ml
Amino acid content [g]	38	48	72	96
Nitrogen content [g]	5.4	6.8	10.2	13.6
Carbohydrate content [g]	120	150	225	300
Lipid content [g]	40	50	75	100

	in 1000 ml	in 1250 ml	in 1875 ml	in 2500 ml
Sodium	40	50	75	100
Potassium	28	35	52.5	70
Magnesium	3.2	4.0	6.0	8.0
Calcium	3.2	4.0	6.0	8.0
Zinc	0.024	0.03	0.045	0.06
Chloride	36	45	67.5	90
Acetate	36	45	67.5	90
Phosphate	12	15	22.5	30

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Emulsion for infusion

Amino acids and glucose solutions: clear, colourless up to straw-coloured solutions

Fat emulsion: oil-in-water emulsion, milky white

	in 1000 ml	in 1250 ml	in 1875 ml	in 2500 ml
Energy in the form of lipids [kJ (kcal)]	1590 (380)	1990 (475)	2985 (715)	3980 (950)
Energy in the form of carbohydrates [kJ (kcal)]	2010 (480)	2510 (600)	3765 (900)	5020 (1200)
Energy in the form of amino acids [kJ (kcal)]	635 (150)	800 (190)	1200 (285)	1600 (380)
Non-protein energy [kJ (kcal)]	3600 (860)	4500 (1075)	6750 (1615)	9000 (2155)
Total energy [kJ (kcal)]	4235 (1010)	5300 (1265)	7950 (1900)	10600 (2530)
Osmolality [mOsm/kg]	1540	1540	1540	1540
Theoretical osmolality [mOsm/l]	1215	1215	1215	1215
pH	5.0 - 6.0	5.0 - 6.0	5.0 - 6.0	5.0 - 6.0

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Supply of energy, essential fatty acids, amino acids, electrolytes and fluids for parenteral nutrition of patients in states of moderately severe catabolism when oral or enteral nutrition is impossible, insufficient or contraindicated. Nutriflex Lipid plus is indicated in adults, adolescents and children older than two years.

### 4.2 Posology and method of administration

#### Posology

The dosage should be adapted to the patients' individual requirements. It is recommended that Nutriflex Lipid plus be administered continuously. A stepwise increase of the infusion rate over the first 30 minutes up to the desired infusion rate avoids possible complications.

#### Adolescents from 14 years of age and adults

The maximum daily dose amounts to 40 ml/kg body weight, corresponding to:

- 1.54 g amino acids /kg body weight per day
- 4.8 g glucose /kg body weight per day
- 1.6 g lipid /kg body weight per day.

The maximum rate of infusion is 2.0 ml/kg body weight per hour, corresponding to:

- 0.08 g amino acids /kg body weight per hour
- 0.24 g glucose /kg body weight per hour
- 0.08 g lipid /kg body weight per hour.

For a patient weighing 70 kg this corresponds to a maximum infusion rate of 140 ml per hour. The amount of substrate administered is then 5.4 g of amino acids per hour, 16.8 g of glucose per hour and 5.6 g of lipids per hour.

#### Paediatric population

#### Newborn infants, infants and toddlers less than two years of age

Nutriflex Lipid plus is contraindicated in newborn infants, infants and toddlers < 2 years of age (see section 4.3).

#### Children from 2 to 13 years of age

The given dosage recommendations are guiding data based on average requirements. The dosage should be individually adapted according to age, developmental stage and illness. For calculation of dosage account must be taken of the hydration status of the paediatric patient.

For children, it might be necessary to start the nutritional therapy with half of the target dosage. The dosage should be increased stepwise according to the individual metabolic capacity up to the maximum dosage.

Daily dose for 2 - 4 years of age: 40 ml/kg body weight, corresponding to:

- 1.54 g amino acids /kg body weight per day
- 4.8 g glucose /kg body weight per day
- 1.6 g lipid /kg body weight per day.

Daily dose for 5 - 13 years of age: 25 ml/kg body weight, corresponding to:

- 0.96 g amino acids /kg body weight per day
- 3.0 g glucose /kg body weight per day
- 1.0 g lipid /kg body weight per day.

The maximum rate of infusion is 2.0 ml/kg body weight per hour, corresponding to:

- 0.08 g amino acids /kg body weight per hour
- 0.24 g glucose /kg body weight per hour
- 0.08 g lipid /kg body weight per hour.

Due to the individual needs of paediatric patients, Nutriflex Lipid plus may not cover sufficiently the total energy and fluid requirements. In such cases carbohydrates and/or lipids and/or fluids must be provided in addition, as appropriate.

#### Patients with renal/hepatic impairment

The doses should be adjusted individually in patients with hepatic or renal insufficiency (see also section 4.4).

#### Duration of treatment

The duration of treatment for the indications stated is not limited. During the administration of Nutriflex Lipid plus it is necessary to provide an appropriate amount of trace elements and vitamins.

#### Duration of infusion of one single bag

The recommended duration of infusion for a parenteral nutrition bag is maximum 24 h.

#### Method of administration

Intravenous use. For central venous infusion only.

### 4.3 Contraindications

- Hypersensitivity to the active substances, to egg, peanut or soya protein or to any of the excipients listed in section 6.1

- Inborn errors of amino acid metabolism

- Severe hypertriglyceridaemia ( $\geq 1000$  mg/dl or 11.4 mmol/l)

- Severe coagulopathy

- Hyperglycaemia not responding to insulin doses of up to 6 units insulin/hour

- Acidosis

- Intrahepatic cholestasis

- Severe hepatic insufficiency

- Severe renal insufficiency in absence of renal replacement therapy

- Aggravating haemorrhagic diatheses

- Acute thromboembolic events, lipid embolism

On account of its composition, Nutriflex Lipid plus must not be used in newborn infants, infants and toddlers under 2 years of age.

General contraindications to parenteral nutrition include:

- Unstable circulatory status with vital threat (states of collapse and shock)

- Acute phases of cardiac infarction and stroke

- Unstable metabolic condition (e.g. severe postaggression syndrome, coma of unknown origin)

- Inadequate cellular oxygen supply

- Disturbances of the electrolyte and fluid balance

- Acute pulmonary oedema

- Decompensated cardiac insufficiency

### 4.4 Special warnings and precautions for use

Caution should be exercised in cases of increased serum osmolality. Disturbances of the fluid, electrolyte or acid-base balance must be corrected before the start of infusion.

Too rapid infusion can lead to fluid overload with pathological serum electrolyte concentrations, hyperhydration and pulmonary oedema.

Any sign or symptom of anaphylactic reaction (such as fever, shivering, rash or dyspnoea) should lead to immediate interruption of the infusion. The serum triglyceride concentration should be monitored when infusing Nutriflex Lipid plus.

Depending on the patient's metabolic condition, occasional hypertriglyceridaemia may occur. If the plasma triglyceride concentration exceeds 4.6 mmol/l (400 mg/dl) during administration of lipids it is recommended to reduce the infusion rate. The infusion must be interrupted if the plasma triglyceride concentration exceeds 11.4 mmol/l (1000 mg/dl), as these levels have been associated with acute pancreatitis.

#### Patients with impaired lipid metabolism

Nutriflex Lipid plus should be administered cautiously to patients with disturbances of lipid metabolism with increased serum triglycerides, e.g. renal insufficiency, diabetes mellitus, pancreatitis, impaired hepatic function, hypothyroidism (with hypertriglyceridaemia), sepsis and metabolic syndrome. If Nutriflex Lipid plus is given to patients with these conditions, more frequent monitoring of serum triglycerides is necessary to assure triglyceride elimination and stable triglyceride levels below 11.4 mmol/l (1000 mg/dl).

In combined hyperlipidaemia and in metabolic syndrome, triglyceride levels react to glucose, lipids and in overnutrition. Adjust dose accordingly. Assess and monitor other lipid and glucose sources, and drugs interfering with their metabolism.

The presence of hypertriglyceridaemia 12 hours after lipid administration also indicates a disturbance of lipid metabolism.

Like all solutions containing carbohydrates, the administration of Nutriflex Lipid plus can lead to hyperglycaemia. The blood glucose level should be monitored. If there is hyperglycaemia, the rate of infusion should be

reduced or insulin should be administered. If the patient is receiving other intravenous glucose solutions concurrently, the amount of additionally administered glucose has to be taken into account.

An interruption of administration of the emulsion may be indicated if the blood glucose concentration rises to above 14 mmol/l (250 mg/dl) during administration.

Refeeding or repletion of malnourished or depleted patients may cause hypokalaemia, hypophosphataemia and hypomagnesaemia. Adequate supplementation of electrolytes according to deviations from normal values is necessary.

Controls of the serum electrolytes, the water balance, the acid-base balance, and of blood cell counts, coagulation status, hepatic and renal function are necessary.

Substitution of electrolytes, vitamins and trace elements may be necessary as required. As Nutriflex Lipid plus contains zinc, magnesium, calcium and phosphate, care should be taken when it is co-administered with solutions containing these substances.

Nutriflex Lipid plus should not be given simultaneously with blood in the same infusion set due to the risk of pseudoagglutination (see also section 4.5).

Nutriflex Lipid plus is a preparation of complex composition. It is, therefore, strongly advisable not to add other solutions (as long as compatibility is not proven - see section 6.2).

As with all intravenous solutions, especially for parenteral nutrition, strict aseptic precautions are necessary for the infusion of Nutriflex Lipid plus.

#### Elderly patients

Basically the same dosage as for adults applies, but caution should be exercised in patients suffering from further diseases like cardiac insufficiency or renal insufficiency that may frequently be associated with advanced age.

#### Patients with diabetes mellitus, impaired cardiac or renal function

Like all large-volume infusion solutions, Nutriflex Lipid plus should be administered with caution to patients with impaired cardiac or renal function.

There is only limited experience of its use in patients with diabetes mellitus or renal failure.

This medicinal product contains 1150 mg sodium per 1250 ml bag, equivalent to 58% of the WHO recommended maximum daily intake of 2 g sodium for an adult.

The maximum daily dose of this product for a 70 kg adult is equivalent to 129% of the WHO recommended maximum daily intake for sodium.

Nutriflex Lipid plus is considered high in sodium. This should be particularly taken into account for those on a low salt diet.

#### Interference with laboratory tests

The fat content may interfere with certain laboratory measurements (e.g. bilirubin, lactate dehydrogenase, oxygen saturation) if blood is sampled before fat has been adequately cleared from the blood stream.

### 4.5 Interaction with other medicinal products and other forms of interaction

Some drugs, like insulin, may interfere with the body's lipase system. This kind of interaction seems, however, to be of only limited clinical importance.

Heparin given in clinical doses causes a transient release of lipoprotein lipase into the circulation. This may result initially in increased plasma lipolysis followed by a transient decrease in triglyceride clearance.

Soya-bean oil has a natural content of vitamin K<sub>1</sub>. This may interfere with the therapeutic effect of coumarin derivatives, which should be closely monitored in patients treated with such drugs.

Potassium-containing solutions like Nutriflex Lipid plus should be used with caution in patients receiving drugs that increase serum potassium concentration, such as potassium-sparing diuretics (triamterene, amiloride, spironolactone), ACE inhibitors (e.g. captopril, enalapril), angiotensin-II-receptor antagonists (e.g. losartan, valsartan), ciclosporin and tacrolimus.

Corticosteroids and ACTH are associated with sodium and fluid retention. Nutriflex Lipid plus should not be given simultaneously with blood in the same infusion set due to the risk of pseudoagglutination (see also section 4.4).

### 4.6 Fertility, pregnancy and lactation

#### Pregnancy

There are no or limited amount of data from the use of Nutriflex Lipid plus in pregnant women. Animal studies are insufficient with respect to reproductive toxicity (see section 5.3). Parenteral nutrition may become necessary during pregnancy. Nutriflex Lipid plus should only be given to pregnant women after careful consideration.

#### Breast-feeding

Components/metabolites of Nutriflex Lipid plus are excreted in human milk, but at therapeutic doses no effects on the breastfed newborns/infants are anticipated.

Nevertheless, breast-feeding is not recommended for mothers on parenteral nutrition.

#### Fertility

No data from the use of Nutriflex Lipid plus available.

### 4.7 Effects on ability to drive and use machines

Nutriflex Lipid plus has no or negligible influence on the ability to drive and use machines.

### 4.8 Undesirable effects

Under conditions of correct use, in terms of dosing monitoring, observation of safety restrictions and instructions, undesirable effects may still occur. The following listing includes a number of systemic reactions that may be associated with the use of Nutriflex Lipid plus.

Undesirable effects are listed according to their frequencies as follows:

Very common ( $\geq 1/10$ )

Common ( $\geq 1/100$  to  $< 1/10$ )

Uncommon ( $\geq 1/1,000$  to  $< 1/100$ )

Rare ( $\geq 1/10,000$  to  $< 1/1,000$ )

Very rare ( $< 1/10,000$ )

Not known (frequency cannot be estimated from the available data)

#### Blood and lymphatic system disorders

Rare: Hypercoagulation

Not known: Leucopenia, thrombocytopenia

#### Immune system disorders

Rare: Allergic reactions (e.g. anaphylactic reactions, dermal eruptions, laryngeal, oral and facial oedema)

#### Metabolism and nutrition disorders

Uncommon: Loss of appetite

Very rare: Hyperlipidaemia, hyperglycaemia, metabolic acidosis

The frequency of these undesirable effects is dose-dependent and may be higher under the condition of absolute or relative lipid overdose.

#### Nervous system disorders

Rare: Headache, drowsiness

#### Vascular disorders

Rare: Hypertension or hypotension, flush

#### Respiratory, thoracic and mediastinal disorders

Rare: Dyspnoea, cyanosis

#### Gastrointestinal disorders

Uncommon: Nausea, vomiting,

#### Hepatobiliary disorders

Not known: Cholestasis

#### Skin and subcutaneous tissue disorders

Rare: Erythema, sweating

#### Musculoskeletal and connective tissue disorders

Rare: Pain in the back, bones, chest and lumbar region

## 5.2 Pharmacokinetic properties

### Absorption

Nutriflex Lipid plus is infused intravenously. Hence, all substrates are available for metabolism immediately.

### Distribution

The dose, rate of infusion, metabolic situation and individual factors of the patient (level of fasting) are of decisive importance for the maximum triglyceride concentrations reached. When used according to the instructions with due regard to the dosage guidelines the triglyceride concentrations do not, in general, exceed 4.6 mmol/l (400 mg/dl). Medium-chain fatty acids have a low affinity to albumin. In animal experiments administering pure medium-chain triglyceride emulsions, it has been shown that medium-chain fatty acids can cross the blood-brain barrier, if overdosed. No adverse effects were observed with an emulsion providing a mixture of medium-chain triglycerides and long-chain triglycerides, as long-chain triglycerides have an inhibiting effect on medium-chain triglyceride hydrolysis. Therefore, toxic effects on the brain can be excluded after the administration of Nutriflex Lipid plus.

Amino acids are incorporated in a variety of proteins in different organs of the body. In addition each amino acid is maintained as free amino acid in the blood and inside cells.

As glucose is water-soluble, it is distributed with the blood over the whole body. At first, the glucose solution is distributed in the intravascular space and then it is taken up into the intracellular space.

No data are available concerning transport of the components through the placental barrier.

### Biotransformation

Amino acids that do not enter protein synthesis are metabolised as follows. The amino group is separated from the carbon skeleton by transamination. The carbon chain is either oxidised directly to CO<sub>2</sub> or utilised as substrate for gluconeogenesis in the liver. The amino group is also metabolised in the liver to urea.

Glucose is metabolised to CO<sub>2</sub> and H<sub>2</sub>O via the known metabolic routes. Some glucose is utilised for lipid synthesis.

After infusion triglycerides are hydrolysed to glycerol and fatty acids. Both are incorporated in physiological pathways for energy production, synthesis of biological active molecules, gluconeogenesis and resynthesis of lipids.

### Elimination

Only minor amounts of amino acids are excreted unchanged in urine.

Excess glucose is excreted in urine only if the renal threshold of glucose is reached.

Both the triglycerides of soya-bean oil and medium-chain triglycerides are completely metabolised to CO<sub>2</sub> and H<sub>2</sub>O. Small amounts of lipids are lost only during sloughing of cells from skin and other epithelial membranes. Renal excretion does virtually not occur.

### 5.3 Preclinical safety data

Non-clinical studies have not been performed with Nutriflex Lipid plus. Toxic effects of mixtures of nutrients given as substitution therapy at the recommended dosage are not to be expected.

### Reproductive toxicity

Phytoestrogens such as β-sitosterol can be found in various vegetable oils, especially in soya-bean oil. Impairment of fertility was determined in rats and rabbits after subcutaneous and intravaginal administration of β-sitosterol. According to the current state of knowledge the observed effects in animals do not seem to have relevance for clinical use.

## 6. PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

Citric acid monohydrate (for pH adjustment)

Glycerol

Egg lecithin

Sodium oleate

Water for injections

### 6.2 Incompatibilities

Nutriflex Lipid plus must not be mixed with other medicinal products for which compatibility has not been documented. See section 6.6.

Nutriflex Lipid plus should not be given simultaneously with blood, see sections 4.4 and 4.5.

### 6.3 Shelf life

#### Unopened

2 years

### After removing the protective overwrap and after mixing of contents of the bag

Chemical and physicochemical in-use stability of the mixture of amino acids, glucose and fat was demonstrated for 7 days at 2-8 °C and additional 2 days at 25 °C.

### After admixture of compatible additives

From a microbiological point of view, the product should be used immediately after admixture of additives. If not used immediately after admixture of additives, in-use storage times and conditions prior to use are the responsibility of the user.

### After first opening (spiking of the infusion port)

The emulsion is to be used immediately after opening of the container.

### 6.4 Special precautions for storage

Do not store above 25 °C.

Do not freeze. If accidentally frozen, discard the bag.

Keep the bag in the outer carton in order to protect from light.

### 6.5 Nature and contents of container

Nutriflex Lipid plus is supplied in flexible multichamber bags of polyamide/polypropylene containing:

- 1250 ml (500 ml of amino acids solution + 250 ml of fat emulsion + 500 ml of glucose solution)
- 1875 ml (750 ml of amino acids solution + 375 ml of fat emulsion + 750 ml of glucose solution)
- 2500 ml (1000 ml of amino acids solution + 500 ml of fat emulsion + 1000 ml of glucose solution)

The multichamber bag is packed in a protective overwrap. An oxygen absorber is placed between the bag and the overwrap; the sachet of inert material contains powdered iron.

The upper left-hand chamber contains a glucose solution, the upper right-hand chamber contains a fat emulsion, and the lower chamber contains an amino acid solution.

The two upper chambers can be connected with the lower chamber by opening the intermediate seam (peel seam).

The design of the bag permits mixing of the amino acids, glucose, lipids and electrolytes in a single chamber. Opening the peel seam results in sterile mixing to form an emulsion.

The different container sizes are presented in cartons containing five bags. Pack sizes: 5 x 1250 ml, 5 x 1875 ml and 5 x 2500 ml

Not all pack sizes may be marketed.

### 6.6 Special precautions for disposal and other handling

No special requirements for disposal.

Parenteral nutrition products should be visually inspected for damage, discolouration and emulsion instability before use.

Do not use bags which are damaged. Overwrap, primary bag and the peel seams between the chambers should be intact. Only use if the amino acid and glucose solutions are clear and colourless up to straw coloured and the lipid emulsion is homogenous with milky white appearance. Do not use if the solutions contain particulate matter. After mixing the three chambers, do not use if the emulsion shows discolouration or signs of phase separation (oil drops, oil layer). Stop the infusion immediately in case of discolouration of the emulsion or signs of phase separation.

#### Preparation of the mixed emulsion:

Remove inner bag from its protective overwrap and proceed as follows:

- Put the bag on a solid, flat surface
- Mix glucose with amino acids by pressing the upper left chamber against the peel seam, then add the fat emulsion by pressing the upper right chamber against the peel seam
- Mix the contents of the bag thoroughly

The mixture is a milky white homogenous oil-in-water emulsion.

#### Preparation for infusion:

The emulsion should always be brought to room temperature prior to infusion.

- Fold the bag and hang it on the infusion stand by the centre hanging loop
- Remove the protective cap from the infusion port and carry out infusion using the standard technique

For single use only.

Container and unused residues must be discarded after use.

Do not reconnect partially used containers.

If filters are used they must be lipid-permeable (pore size ≥ 1.2 µm).

## 7. DATE OF REVISION OF THE TEXT

02/2022

**B | BRAUN**

B. Braun Melsungen AG  
34209 Melsungen  
Germany

